

Talking to your loved one's healthcare provider about LONSURF[®] (trifluridine and tipiracil) tablets

For those with colon or rectal cancer that has spread to other parts of the body and who have previously been treated with or cannot receive certain chemotherapy medicines, LONSURF might be another treatment option.

Your loved one's doctor or nurse can help you find out whether LONSURF is right for your loved one. Be sure to bring this guide to your loved one's next appointment, and use the questions on the following pages to help start the conversation.

LONSURF may be an option for patients who have been previously treated with or cannot receive some or all of the following treatments:

- **Fluoropyrimidine** - such as 5-fluorouracil, also known as 5-FU
- **Oxaliplatin-based treatment** - such as FOLFOX (5-FU, leucovorin, and oxaliplatin); CAPOX (capecitabine and oxaliplatin); or Eloxatin[®] (oxaliplatin)
- **Irinotecan-based treatment** - such as FOLFIRI (5-FU, leucovorin, and irinotecan); CAPIRI (capecitabine and irinotecan); or Camptosar[®] (irinotecan)
- **Anti-VEGF biological therapy, or VEGF inhibitors** - such as Avastin[®] (bevacizumab); or Zaltrap[®] (ziv-aflibercept)
- **Anti-EGFR therapy, or EGFR inhibitors** (if you have the **KRAS wild type gene**) - such as Erbitux[®] (cetuximab); or Vectibix[®] (panitumumab)

Registered trademarks are the property of their respective owners.

Here are some questions to ask your loved one's healthcare provider at their next appointment:

1. Based on the treatments my loved one has previously taken, is he/she a candidate for LONSURF? _____

2. How can LONSURF help? _____

3. How and where is LONSURF taken? _____

4. What dose of LONSURF would he or she take? How often? _____

5. How often will we need to come into the office with this treatment? _____

6. What happens if a dose is missed? _____

7. What side effects could my loved one have? _____

Doctor Discussion Guide



8. When should I call you if he or she is having signs and symptoms of side effects? _____

9. How can I help my loved one manage any side effects? _____

10. How will the treatment affect my loved one's daily activities or ability to work? _____

11. Does Taiho Oncology offer any financial assistance programs? _____

INDICATION

LONSURF® (trifluridine and tipiracil) tablets are a prescription medicine used to treat people with colon or rectal cancer that has spread to other parts of the body and who have been previously treated with or cannot receive certain chemotherapy medicines. It is not known if LONSURF is safe and effective in children.

IMPORTANT SAFETY INFORMATION

LONSURF may cause serious side effects, including:

- **Low blood counts.** Low blood counts are common with LONSURF and can sometimes be severe and life-threatening. LONSURF can cause a decrease in your white blood cells, red blood cells, and platelets. Low white blood cells can make you more likely to get serious infections that could lead to death. Your healthcare provider should do blood tests before you receive LONSURF, at day 15 during treatment with LONSURF, and as needed to check your blood cell counts. Your healthcare provider may lower your dose of LONSURF or stop LONSURF if you have low white blood cell or platelet counts

Tell your healthcare provider right away if you get any of the following signs and symptoms of infection during treatment with LONSURF: fever, chills, or body aches.

Before taking LONSURF, tell your healthcare provider about all of your medical conditions, including if you:

- Have kidney or liver problems
- Are pregnant or plan to become pregnant. LONSURF can harm your unborn baby
 - **Females** who can become pregnant should use effective birth control during treatment with LONSURF. Tell your healthcare provider immediately if you become pregnant
 - **Males**, while on treatment and for 3 months after your last dose of LONSURF, you should use a condom during sex with female partners who are able to become pregnant. Tell your healthcare provider right away if your partner becomes pregnant while you are taking LONSURF
- Are breast-feeding or plan to breast-feed. It is not known if LONSURF passes into your breast milk. Do not breast-feed during treatment with LONSURF and for 1 day after your last dose of LONSURF

Tell your healthcare provider about all the prescription and over-the-counter medicines, vitamins, and herbal supplements you take.

The **most common side effects** with LONSURF include tiredness, nausea, decreased appetite, diarrhea, vomiting, abdominal pain, and fever.

Tell your doctor if you have nausea, vomiting, or diarrhea that is severe or that does not go away.

These are not all of the possible side effects of LONSURF. For more information, ask your healthcare provider. Call your doctor for medical advice about side effects.

Please see the full Prescribing Information at LONSURF.com.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.